Impairment of proteostasis network in Down syndrome prior to the development of Alzheimer's disease neuropathology: Redox proteomics analysis of human brain

Fabio Di Domenico [a], Raffaella Coccia[a], Annalisa Cocciolo [a], M. Paul Murphy [d], Giovanna Cenini [b,c], Elizabeth Head [c,d], D. Allan Butter eld [b,d,e], Alessandra Giorgi [a], Maria Eugenia Schinina [a], Cesare Mancuso[f], Chiara Cini [a], Marzia Perluigi [a,]

[a] Department of Biochemical Sciences, Sapienza University of Rome, 00185 Rome, Italy
[b] Department of Chemistry, University of Kentucky, Lexington, KY 40506-0055, USA
[c] Department of Molecular and Biomedical Pharmacology, University of Kentucky, Lexington, KY 40506-0055, USA
[d] Sanders-Brown Center on Aging, University of Kentucky, Lexington, KY 40506-0055, USA
[e] Center of Membrane Sciences, University of Kentucky, Lexington, KY 40506-0055, USA
[f] Institute of Pharmacology, Catholic University School of Medicine, Largo F. Vito, 1, 00168 Rome, Italy

a r t i c l e i n f o

Article history:
Received 26 November 2012
Received in revised form 18 March 2013
Accepted 10 April 2013
Available online 18 April 2013

Keywords:
Down syndrome
Oxidative stress
Autophagy
Proteasome
Alzheimer disease
Trisomy 21

a b s t r a c t

DS is the most frequent genetic cause of intellectual disability characterized by the anomalous presence of three copies of chromosome 21. One of the peculiar features of DS is the onset of Alzheimer's disease neuropathology after the age of 40 years characterized by deposition of senile plaques and neuro brillary tangles. Growing studies demonstrated that increased oxidative damage, accumulation of unfolded/damaged protein aggregates and dysfunction of intracellular degradative system are key players in neurodegenerative processes. In this study, redox proteomics approach was used to analyze the frontal cortex from DS subjects under the age of 40 compared with age-matched controls, and proteins found to be increasingly carbonylated were identi ed. Interestingly, our results showed that oxidative damage targets speci cally different components of the intracellular quality control system such as GRP78, UCH-L1, V0-ATPase, cathepsin D and GFAP that couples with decreased activity of the proteasome and autophagosome formation observed. We also reported a slight but consistent increase of A 1..42 SDS- and PBS-soluble form and tau phosphorylation in DS versus CTR. We suggest that disturbance in the proteostasis network could contribute to the accumulation of protein aggregates, such as amyloid deposits and NFTs, which occur very early in DS. It is likely that a sub-optim [(1.)-302(Introd)11(uction)]TJ /T1_0 1 Tf 1.501 -2.625 Td [(Do)25(w)15(n)-214(sy)25(nd)23(r)15(o)14(me)-200

1% (w/v) sodium dodecyl sulfate (SDS), 30% (v/v) glycerol, and 0.5% DTT and again for another 10 min in the same buffer containing 4.5% iodoacetamide in place of DTT. The second dimension was performed using 12% precast Criterion gels (Bio-Rad). The gels were incubated in xing solution (7% acetic acid, 10% methanol) for 20 min and then stained for 1 h in Bio-Safe Coomassie gel stain (Bio-Rad, Hercules, CA, USA) and destained overnight in deionized water. The Coomassie gels were scanned using a GS 800 densitometer (Bio-Rad, Hercules, CA, USA).

2.4. 2D oxyblot

For 2D OxyBlot, 2D gels (200 g of proteins) were blotted onto nitrocellulose membranes (Bio-Rad, Hercules, CA, USA) and 2,4-dinitrophenylhydrazine (DNPH) derivatization was performed. Brie2,4-

2.7. Western blot

For Western blot, 30 g of proteins (CTR and DS) were separated by 12% SDS..PAGE and blotted onto a nitrocellulose membrane (Bio-Rad, Hercules, CA, USA). Membranes were blocked with 3% bovine serum albumin in T-TBS and incubated for 1 h and 30 min at room temperature with primary anti-LC3 antibody (1:100; Novus Biologicals, Littleton, CO, USA), anti proteasome 20S core subunit (1:1000; ENZO Life Science, Farmingdale, NY, USA) and 19S 10B (1:500; Novus Biologicals, Littleton, CO, USA), anti-e, Farmin CO, No00;

3.4. Cathepsin D and UCH-L1 activity

To analyze if the increased carbonylation of CatD may affect protein function, we measured CatD enzyme activity in the two groups of brain samples. A signi cant ($p < 0.05$) slight increase of CatD activity in DS samples, of about 10% compared to CTR samples, was demonstrated (Fig. 5A). As well, the activity of UCH-L1 was measured in order to determine any alteration on its functionality.

Interestingly, UCH-L1 enzyme activity present a decrease of about 30% in DS compared to CTR.

3.5. LC3 II/I levels

As the first step, the protective mechanism used by cells to adapt to stress of the ER is the induction of members of the glucose-regulated protein (Grp) family, mainly GRP78. The induction of mammalian Grp proteins in response to ER stress involves a complex network of regulators and novel mechanisms. Our data demonstrate a relationship between OS and protein misfolding in DS brain, showing that GRP78 can be a target of OS in vivo. Oxidative damage may result in GRP78 dysfunction (inability to bind to the misfolded proteins), which provides a conceivable mechanistic link between deficits in molecular chaperones, accumulation of misfolded proteins, and risk of cognitive decline. According to our previous studies [20,37], increased levels of GRP78 indicate the induction of the UPS system, as results of the presence of unfolded/misfolded proteins; however, the pro-oxidant environment in DS appears to result in GRP78 oxidation that might lead to dysfunction of the UPR system.

Upon activation of ER-stress response, ERAD mediates ubiquitination

Fig. 4. Panel A: Bar graph of proteasome subunits (20S and 19S) levels in CTR and DS samples measured by WB analysis. Panel B: Proteasome functionality measured by enzymatic assay of chymotrypsin-like, trypsin-like and caspase-like activity in CTR and DS samples. *p < 0.01.

laboratory and others demonstrated that the levels of UCH-L1 are decreased in AD brain and also that this enzyme is oxidatively modified [44]. One of the major consequences of aberrant UCH-L1 activity is an impaired proteasome proteolytic system, which will lead to accumulation of damaged proteins and formation of protein aggregates. The proteolytic core of the proteasome or •20S proteasome Ž is involved in the degradation of oxidatively modified proteins [46..48]. Oxidatively modified proteins can undergo chemical fragmentation or form aggregates because of covalent cross-linking reactions and increased surface hydrophobicity [49]. The recognition of hydrophobic amino acid residues of oxidized proteins and their subsequent degradation by the 20S proteasome could be a selective mechanism to remove oxidatively damaged proteins from the cell [50]. Although we found increased levels of the 20S subunit, the activities of trypsin-like, chymotrypsin-like and caspase-like of the proteasome were decreased in DS brain compared with age-matched controls. It is likely that proteasome activation may be a signal to enhance protein degradation in affected neurons to prevent the formation of toxic aggregates, but at the same time the enzyme is directly or indirectly damaged by ROS and its functionality is affected.

Another major degradation system reported to be activated by ER stress in vitro is autophagy [51..53]. During autophagy, a double membrane is wrapped around an organelle or protein aggregate and this autophagosome subsequently fuses with a lysosome, forming an autophagolysosome where degradation takes place. Healthy neurons only rarely show autophagic structures, whereas many autophagic vacuoles can be observed in AD neuronal cell bodies and dystrophic neuritis [54].

Autophagy is considered as a basal cellular mechanism to control homeostasis of intracellular proteins and organelles. However, autophagy can be also induced to protect cells under various physiological stresses such as nutrient depletion, genotoxic agents, aggregated proteins, or cytokines. In particular, if the first defense mechanism by non-enzymatic molecules and enzymatic scavengers is overwhelmed under oxidative stress, autophagy plays a crucial role in cell protection



Fig. 5. Panel A: Cathepsin D activity in CTR ad DS samples measured by fluorescent assay at 328/393 Ex/Em. Panel B: UCH-L1 activity assay in CTR and DS samples measured by fluorescent assay at 380/460 Ex/Em n = 8; *p < 0.05.

from harmful oxidized materials by effectively removing them [55]. Unfortunately, these defense mechanisms are not always enough to overcome the oxidative stress, resulting in autophagic cell death (type II programmed cell death) that is distinct from apoptosis (type I programmed cell death) [56]. However, the mechanistic basis coordinating autophagic processes under oxidative stress is still not clear and further studies are needed.

We hypothesize that the oxidation of V_0-ATPase pump could lead to autophagy dysfunction. This proton pump is essential for acidic lysosomal pH. Mutation of lysosomal ATPase genes is a well-recognized risk factor for autophagy related neurodegenerative diseases [57,58]. It is reasonable to speculate that once oxidized V_0-ATPase has altered ability to regulate intracellular pH thus affecting proper lysosome functionality and autophagy, as was previously seen in a PD model [59]. In

[63] S. Grkovic, V.C. O'Reilly, S. Han, M. Hong, R.C. Baxter, S.M. Firth, IGFBP-3 binds GRP78, stimulates autophagy and promotes the survival of breast cancer cells exposed to adverse microenvironments, Oncogene (Jul. 2 2012), http: